



Review

CSF-1 as a Regulator of Macrophage Activation and Immune Responses

MATTHEW J. SWEET* and DAVID A. HUME

CRC for Chronic Inflammatory Diseases, Institute for Molecular Bioscience and Departments of Microbiology/Parasitology and Biochemistry, University of Queensland, Qld 4072, Australia

Abstract. Macrophage activation is a key determinant of susceptibility and pathology in a variety of inflammatory diseases. The extent of macrophage activation is tightly regulated by a number of pro-inflammatory cytokines (e.g. IFN- γ , IL-2, GM-CSF, IL-3) and anti-inflammatory cytokines (e.g. IL-4, IL-10, TGF- β). Macrophage colony-stimulating factor (CSF-1/M-CSF) is a key differentiation, growth and survival factor for monocytes/macrophages and osteoclasts. The role of this factor in regulating macrophage activation is often overlooked. This review will summarize our current understanding of the effects of CSF-1 on the activation state of mature macrophages and its role in regulating immune responses.

Key words: CSF-1; macrophage; lipopolysaccharide; Toll-like receptors; inflammation.

Introduction

The production of circulating monocytes and tissue macrophages from the bone marrow is dependent on colony-stimulating factor 1 (CSF-1). This is highlighted by the gross deficiencies in macrophage development and numbers that occur in the *op/op* mouse, which has a natural inactivating mutation in the CSF-1 gene (for review see⁸⁴), and by the demonstration that administration of CSF-1 to mice caused a 10-fold increase in blood monocyte numbers and increased macrophage numbers in the liver, spleen and peritoneal cavity⁴¹.

Via alternative splicing, post-translational modifications and proteolytic processing, CSF-1 is produced in multiple forms; a secreted homodimeric glycoprotein, a secreted proteoglycan and a membrane-bound glycoprotein^{73, 84}. The CSF-1 receptor (CSF-1R), encoded by the *c-fms* protooncogene, is a type III receptor tyrosine kinase with structural similarity to c-kit and the platelet-derived growth factor receptor and the *fms*-like receptors *flk*, and *flt* 1, 2 and 3. Mice with a targeted disruption of the *c-fms* gene are essentially a phenocopy of the *op/op* mouse, indicating that all of the actions of CSF-1 are mediated by the CSF-1R¹⁸. Ligand binding to the CSF-1R initiates receptor dimerization, auto-

Abbreviations used: BMM – bone marrow-derived macrophage, CpG DNA – bacterial CpG-containing DNA, CSF-1 – colony-stimulating factor 1, CSF-1R – CSF-1 receptor, EBV – Epstein-Barr virus, HIV-1 – human immunodeficiency virus type 1, IDO – indoleamine 2,3-dioxygenase, IL – interleukin, LPS – lipopolysaccharide, MMP – matrix metalloproteinase, RA – rheumatoid arthritis, TLR – Toll-like receptor, TNF – tumor necrosis factor.

* Correspondence to: Dr. Matthew Sweet, Institute for Molecular Bioscience University of Queensland, Qld 4072, Australia, tel.: +61 7 3346 2072, fax: +61 7 3346 2101, e-mail: m.sweet@imb.uq.edu.au

phosphorylation, and activation of multiple signalling cascades including the mitogen-activated protein kinase and P-I-3 kinase/Akt pathways^{21, 35}.

Although relatively few studies have addressed the biological roles of the different CSF-1 isoforms, it is likely that they have distinct functions *in vivo*. For example, the proteoglycan form is localized to specific types of extracellular matrix which might enable specialized functions such as regulation of bone homeostasis⁶⁸. Daily administration of soluble recombinant CSF-1 to the *op/op* mouse from day 3 of life was able to correct some but not all of the deficiencies in macrophage numbers in different tissues¹³. This supports the argument that the full biological activity of CSF-1 requires all isoforms and not just soluble circulating CSF-1, although it is also possible that pre-natal absence of CSF-1 has effects that cannot be subsequently restored by exogenous CSF-1. The gene regulatory elements that are sufficient to reconstitute normal CSF-1 expression *in vivo* have been defined⁷⁷. Transgenic expression on the *op/op* background of cDNAs encoding the secreted, proteoglycan or membrane-bound isoforms under the control of these elements should therefore provide clear evidence for any isoform-specific functions.

Regardless of the functions of the different isoforms, it is clear that CSF-1 is abundantly present *in vivo*. The major source of circulating CSF-1 is thought to be endothelial cells that line blood vessels⁷⁶, but a range of other cell types, including fibroblasts, osteoblasts, monocytes, B cells, T cells and bone marrow stromal cells also produce CSF-1. In mice, CSF-1 levels are in the order of tens of nanograms per millilitre in the circulation and tens of picograms per milligram in a variety of tissues including liver, lung, spleen, kidney, intestine and heart⁷⁵. CSF-1 levels are dramatically increased upon challenge with lipopolysaccharide (LPS)⁷⁵ or with infectious agents such as *Listeria monocytogenes*^{15, 32} and *Candida albicans*¹⁴. In humans, a similar situation is apparent; CSF-1 levels were enhanced in patients with sepsis²⁹, and LPS administration to cancer patients increased CSF-1 levels²³. These observations would suggest that CSF-1 is likely to regulate immune responses to infection in both mice and humans.

Macrophage-Priming by CSF-1

The requirement for the CSF-1/CSF-1R system in macrophage development makes interpretation of *in vivo* studies aimed at addressing the role of CSF-1 in

immunological responses difficult, since modulation of CSF-1 action affects both monocyte/macrophage production and survival, as well as mature macrophage functions. Additionally, CSF-1 is required for the maintenance of protein synthesis. JESSUP et al.⁴³ showed that CSF-1 greatly increased acetylated low-density lipoprotein uptake and foam cell formation, but this was entirely attributable to an increase in cell size; no effect was apparent when these parameters were assessed on a per microgram protein basis. This absolute requirement on CSF-1 for normal macrophage function raises an important issue with regard to the many *in vitro* studies that are performed on this cell type. Since it is normally present *in vivo* and because its absence has gross effects on normal cellular functions, CSF-1 should be present in culture medium during *in vitro* studies on macrophage function. This is often not the case, particularly in immunological studies.

A role for CSF-1 in regulating macrophage activation and inflammation is supported by *in vitro* studies in which CSF-1 enhanced macrophage phagocytic activity against *Listeria monocytogenes*¹⁶, microbicidal action against *Leishmania mexicana*⁴⁰, *Candida albicans*⁴⁷ and *Aspergillus fumigatus*⁷⁴ and primed tumoricidal activity upon triggering with activating agents^{66, 79, 80}. In other studies, CSF-1 did not enhance the ability of macrophages to kill intracellular organisms, including *Leishmania major*⁶, *Mycobacterium tuberculosis*¹⁹ and *Listeria monocytogenes*²⁰. The basis for this inconsistency is unclear.

In contrast to *in vitro* studies, *in vivo* studies strongly support a role for CSF-1 in regulating primary immune responses to infection. For example, mice challenged with *Listeria monocytogenes* had an increased bacterial load following administration of an anti-CSF-1 antibody at the time of challenge³² and CSF-1 transgenic mice were more resistant to *Listeria monocytogenes* than wild-type controls⁸. CSF-1 administration also had protective effects for the host against challenge with *Listeria monocytogenes*⁴⁹ and *Candida albicans*¹⁴.

As well as enhancing the ability of macrophages to destroy invading pathogens directly, CSF-1 is also a potent regulator of monokine production that regulates the magnitude of the inflammatory response. CSF-1 alone triggered production of mRNAs for interleukin (IL)-6, granulocyte macrophage (GM)-CSF, IL-1 α and IL-1 β in murine resident peritoneal macrophages^{25, 45}. In other studies with different macrophage populations, CSF-1 alone did not trigger pro-inflammatory cytokine production, but instead acted as a potent priming signal for a subsequent activation stimulus. Pre-treatment of thioglycollate-elicited peritoneal macrophages or bone

marrow-derived macrophages (BMM) with CSF-1-enhanced production of key pro-inflammatory cytokines, including IL-6, IL-12 and tumor necrosis factor (TNF)- α , upon triggering with LPS^{26, 45, 46, 86}, TNF- α ²⁶ or extracellular matrix proteins^{51, 52}. Similar priming effects of CSF-1 have been reported for human monocytes^{3, 80, 98}. These *in vitro* studies correlate with *in vivo* studies in which the *op/op* mouse was resistant to the toxic effects of LPS^{87, 100}, although the interpretation of this data is somewhat compromised by the defects in macrophage development in the *op/op* mouse.

Interestingly, the ability of CSF-1 to prime macrophage activation is dependent upon the nature of the activating stimulus. Whereas CSF-1 synergized with LPS for pro-inflammatory cytokine production from BMM, it suppressed these responses when the activating signal was bacterial CpG-containing DNA (CpG DNA) and had no effect when the stimulus was bacterial lipopeptide⁸⁶. Hence, the role of CSF-1 may depend upon the type of infectious challenge. In Gram-negative infections where LPS will be detected, CSF-1 may act as a potent priming signal for macrophage activation, whilst in a Gram-positive infection CSF-1 may not be as critical in regulating macrophage function.

The differential effects of CSF-1 on responses to different activating signals is, in part, mediated by selective effects of CSF-1 on expression of Toll-like receptors (TLRs)⁸⁶, which are required for signaling in response to microbial products⁸⁹. Whereas CSF-1 did not regulate expression of a variety of TLR members, including TLR3, 4, 5 and 7, expression of TLR9, which is required for responses to CpG DNA, was down-regulated by CSF-1. Levels of TLR2 and 6 mRNA, encoding receptors for Gram-positive components, were also moderately suppressed by CSF-1 (unpublished data and ⁸⁶).

The biological significance of the potent inhibitory effect of CSF-1 on TLR9 expression and responses to CpG DNA is not clear, but others have also reported that CSF-1 can in some cases inhibit macrophage activation. For example, CSF-1 pretreatment inhibited the macrophage respiratory burst when zymosan, a TLR2/TLR6 agonist⁶⁹, was used as the triggering signal⁷⁰. Hence, CSF-1 may have dual roles in responses to infection; as a positive regulator of macrophage activation when extreme danger signals such as LPS are detected and to prevent excessive macrophage activation when less hazardous products such as CpG DNA and zymosan are encountered.

Mechanisms of CSF-1-Mediated Priming of Cytokine Production

In human monocytes/macrophages, the mechanisms by which CSF-1 primes activation have not been extensively studied. CSF-1 marginally increased expression of CD14 on human monocytes and this may partially account for increased sensitivity to LPS³. Whether CSF-1 regulates LPS recognition in murine macrophages has not been adequately addressed, but CSF-1 did not regulate expression of the receptor for LPS, TLR4 or early signaling events in response to maximal LPS doses⁸⁶. Instead, CSF-1 appears to prime murine macrophage responses to LPS through a number of distinct mechanisms. The synergism between CSF-1 and either LPS or TNF- α for IL-6 production, from resident peritoneal macrophages was largely-mediated by CSF-1-stimulated autocrine GM-CSF production, since synergism was greatly reduced, although not abolished, in GM-CSF-deficient mice²⁶. The mechanism by which CSF-1 augments IL-12 production is not known, but may also involve autocrine GM-CSF, since this factor synergises with LPS for IL-12 production (unpublished data and ³⁹). Another possibility is that synergy occurs at the transcriptional level, since Ets-2, which binds to and activates the IL-12 p40 promoter^{56, 57}, is activated in a sustained fashion in response to CSF-1²⁸. The priming effect for TNF- α production is downstream of mRNA regulation since CSF-1 enhanced LPS-induced TNF- α secretion but not LPS-induced TNF- α mRNA levels in BMM⁸⁶. Since CSF-1 is known to increase production of matrix metalloproteinase (MMP) ⁹², one possibility is that CSF-1 also induces production of the MMP-related protein TNF- α converting enzyme that is required for cleavage of membrane-bound TNF- α ^{60, 61}.

CSF-1 as a Regulator of Leukocyte Recruitment

Apart from affecting macrophage activation, CSF-1 acts at multiple levels to regulate cellular recruitment, a hallmark of inflammatory responses. Firstly, CSF-1 has pronounced effects on macrophage motility. Treatment of the CSF-1-dependent murine macrophage cell line BAC1.2F5 with CSF-1 rapidly triggered actin reorganization, membrane ruffling and cell spreading^{2, 11}. Further, CSF-1 had chemokinetic and chemotactic activity on human monocytes⁹⁶, BAC1.2F5 cells⁹⁹ and myeloid progenitor cells-transfected with a *c-fms* expression plasmid⁷¹. Thus, the enhanced production of CSF-1 during responses to infection may serve to re-

cruit monocytes rapidly to the inflammatory site. Secondly, CSF-1 regulates expression of many genes that encode mediators of adhesion and migration. These include the integrins, $\alpha 4\beta 1$ and $\alpha 5\beta 1$ ⁸², urokinase plasminogen activator^{36, 83}, plasminogen activator inhibitors^{17, 36, 101} and MMP9⁹². Finally, CSF-1 may enhance chemokine production from monocytes/macrophages, thereby allowing for recruitment of other effector cells; CSF-1 increased IL-8 production from human monocytes^{38, 90}. The importance of CSF-1 for the recruitment of monocytes during inflammatory responses has also been documented *in vivo*. Blockade of CSF-1 action suppressed monocyte recruitment during *Listeria monocytogenes* infection³¹ and renal inflammation⁵⁴.

CSF-1 Involvement in Viral Infections

Many viruses, including respiratory syncytial virus, measles virus, dengue virus and HIV-1, replicate within macrophages as a means of escaping immune detection, and in many cases CSF-1 regulates macrophage function during viral infection. This is particularly true for HIV-1, where CSF-1 is an important factor in the HIV-1 replicative strategy. Firstly, HIV-1 replication has been associated with increased CSF-1 production *in vivo*³⁰ and replication of HIV-1 within monocyte-derived macrophages triggered CSF-1 production^{33, 53}. Secondly, CSF-1 enhanced HIV-1 replication in macrophages^{7, 44}, possibly by regulating expression of the HIV-1 co-receptor CCR5⁹⁷. Thus, HIV-1 appears specifically to induce CSF-1 production from macrophages and this, in turn, enhances its ability to replicate within this cell type. CSF-1/CSF-1 receptor antagonists may therefore have therapeutic potential as agents to block HIV-1 replication within macrophages.

Whilst CSF-1 is a positive regulator of HIV-1 replication, it may be protective to the host for other viral pathogens. Intranasal administration of CSF-1 offered almost complete, protection to BALB/c mice upon challenge with otherwise lethal doses of Sendai virus⁵⁹, and CSF-1 treatment of macrophages *in vitro* induced resistance to vesicular stomatitis virus⁵⁵. Interestingly, Epstein-Barr virus (EBV) encodes a protein, BARF1, that bound to and neutralized CSF-1⁸⁵. Although the biological consequences of this have not been addressed, one might predict that blockade of CSF-1 signaling via BARF1 would provide a selective advantage to EBV over the host. Indeed, there are reports of CSF-1R expression on B cells^{4, 91}, which may be relevant to EBV infection. Whilst some of the anti-viral actions of CSF-1 may be attributed to production of type 1 and

type 2 interferons^{55, 59}, one report demonstrated that a CSF-1 expression construct, when administered with a DNA vaccine, enhanced cytotoxic T lymphocyte responses⁵⁰. Therefore, CSF-1 may also enhance the development of antigen-specific anti-viral mechanisms.

Effects of CSF-1 on the Acquired Response

Whilst most of the studies described above indicate that CSF-1 enhances macrophage activation and the innate response, CSF-1 is actually immunosuppressive for antigen-specific responses. T cell responses against allogeneic cells were suppressed in the presence of CSF-1-stimulated macrophages from tumor-bearing hosts⁹⁵, and CSF-1 suppressed MHC class II expression in the placenta⁹⁴. Other studies have substantiated this immunosuppressive effect of CSF-1 *in vitro*^{62, 63}, and CSF-1 administration *in vivo* suppressed proliferation of purified splenocytes to T cell mitogens²².

Apart from targeting MHC class II expression, the immunosuppressive effect of CSF-1 may be dependent on the enzyme indoleamine 2,3-dioxygenase (IDO), which was induced in co-cultures of CSF-1-derived macrophages and T cells⁶⁴. IDO rapidly degrades tryptophan, an amino acid that is essential for T cell proliferation. Several studies have reported inducible expression of membrane-bound CSF-1 on T cells upon activation^{34, 72, 88, 102}. Given the inhibitory effects of CSF-1 on T cell proliferation, inducible expression of CSF-1 in T cells may provide a negative feedback mechanism to arrest T cell proliferation via antigen-presenting cell-mediated tryptophan depletion. Despite these studies, the relevance of CSF-1 as an immunosuppressive agent during antigen-specific responses to infectious agents remains to be established. However, CSF-1-mediated effects on T cell function may be relevant to tolerance of the allogeneic fetus during pregnancy. IDO expression in the placenta is critical for maternal T cell tolerance⁶⁵, and the most dramatic regulation of CSF-1 *in vivo* occurs during pregnancy where uterine, but not circulating, CSF-1 levels increase 1000 fold⁵. Hence, CSF-1-induced IDO may contribute to the maintenance of tolerance to the developing fetus.

CSF-1 and Chronic Inflammatory Responses

The involvement of macrophages in chronic inflammatory diseases such as rheumatoid arthritis (RA) is well established and the effects of CSF-1 in such diseases have also been studied. In collagen-induced arth-

ritis, a mouse model of RA, administration of CSF-1 exacerbated disease severity whilst an anti-CSF-1 antibody reduced the severity of established arthritis¹². CSF-1 has also been implicated as a contributor to disease severity in other arthritic models^{1, 9}. Evidence exists for CSF-1 involvement in RA itself; CSF-1 levels were elevated in RA patient sera⁴⁸ and synovial fluid²⁷, and synovial fibroblasts from RA patients produce CSF-1⁸¹.

Apart from arthritic diseases, there is an extensive literature on the contribution of CSF-1 to kidney disease. Macrophage accumulation is a predictor of renal outcome in glomerulonephritis and correlates with kidney dysfunction in humans⁶⁷, and elevated levels of renal CSF-1 are apparent in glomerulonephritis patients⁴². Other studies have also documented enhanced CSF-1 levels in sera of patients with chronic renal disease^{58, 78}. In experimental disease models, there is clear evidence for the involvement of CSF-1 in directing excessive macrophage proliferation and tissue damage. The severity of lupus nephritis in MRL-lpr mice correlated with CSF-1 levels¹⁰, treatment with anti-CSF-1R antibody reduced local macrophage proliferation during experimentally induced renal inflammation⁵⁴, and treatment of mice with CSF-1 enhanced LPS-induced glomerular macrophage accumulation and proteinuria⁹³.

Conclusion

Determination of the exact roles of CSF-1 during immune responses has been difficult because of its essential role in the differentiation, proliferation and normal cellular function of macrophages. Nonetheless, numerous studies have documented effects of CSF-1 on macrophage activation *in vitro* and inflammatory responses *in vivo*. The fact that CSF-1 levels are elevated during responses to infection in mice and humans strongly supports the concept that this factor has key immunoregulatory functions. As with other pro-inflammatory cytokines, CSF-1 exerts protective effects against many infectious agents (with some notable exceptions), but elevated levels of CSF-1 may contribute to pathology in both acute and chronic inflammatory diseases. Hence, specific antagonists of CSF-1 action may have therapeutic potential as a means of preventing pathology associated with excessive macrophage accumulation and activation.

Acknowledgment. This work was supported by the NHMRC of Australia through a C. J. Martin fellowship (Reg Key No. 987038).

References

1. ABD A. H., SAVAGE N. W., HALLIDAY W. J. and HUME D. A. (1991): The role of macrophages in experimental arthritis induced by *Streptococcus agalactiae* sonicate: actions of macrophage colony-stimulating factor (CSF-1) and other macrophage-modulating agents. *Lymphokine Cytokine Res.*, **10**, 43–50.
2. ALLEN W. E., JONES G. E., POLLARD J. W. and RIDLEY A. J. (1997): Rho, Rac and Cdc42 regulate actin organization and cell adhesion in macrophages. *J. Cell Sci.*, **110**, 707–720.
3. ASAKURA E., HANAMURA T., UMEMURA A., YADA K., YAMAUCHI T. and TANABE T. (1996): Effects of macrophage colony-stimulating factor (M-CSF) on lipopolysaccharide (LPS)-induced mediator production from monocytes *in vitro*. *Immunobiology*, **195**, 300–313.
4. BAKER A. H., RIDGE S. A., HOY T., CACHIA P. G., CULLIGAN D., BAINES P., WHITTAKER J. A., JACOBS A. and PADUA R. A. (1993): Expression of the colony-stimulating factor 1 receptor in B lymphocytes. *Oncogene*, **8**, 371–378.
5. BARTOCCI A., POLLARD J. W. and STANLEY E. R. (1986): Regulation of colony-stimulating factor 1 during pregnancy. *J. Exp. Med.*, **164**, 956–961.
6. BELOSEVIC M., DAVIS C. E., MELTZER M. S. and NACY C. A. (1988): Regulation of activated macrophage antimicrobial activities. Identification of lymphokines that cooperate with IFN- γ for induction of resistance to infection. *J. Immunol.*, **141**, 890–896.
7. BERGAMINI A., PERNO C. F., DINI L., CAPOZZI M., PESCE C. D., VENTURA L., CAPPANNOLI L., FALASCA L., MILANESE G., CALIO R. and ROCCHI G. (1994): Macrophage colony-stimulating factor enhances the susceptibility of macrophages to infection by human immunodeficiency virus and reduces the activity of compounds that inhibit virus binding. *Blood*, **84**, 3405–3412.
8. BERNIER T., HALTER R., PAU D., RITTINGHAUSEN S. and EM-MENDORFFER A. (2001): M-CSF transgenic mice: role of M-CSF in infection and autoimmunity. *Exp. Toxicol. Pathol.*, **53**, 165–173.
9. BISCHOF R. J., ZAFIROPOULOS D., HAMILTON J. A. and CAMPBELL I. K. (2000): Exacerbation of acute inflammatory arthritis by the colony-stimulating factors (CSF)-1 and granulocyte macrophage (GM)-CSF: evidence of macrophage infiltration and local proliferation. *Clin. Exp. Immunol.*, **119**, 361–367.
10. BLOOM R. D., FLORQUIN S., SINGER G. G., BRENNAN D. C. and KELLEY V. R. (1993): Colony-stimulating factor-1 in the induction of lupus nephritis. *Kidney Int.*, **43**, 1000–1009.
11. BOOCOCK C. A., JONES G. E., STANLEY E. R. and POLLARD J. W. (1989): Colony-stimulating factor-1 induces rapid behavioural responses in the mouse macrophage cell line, BAC1.2F5. *J. Cell Sci.*, **93**, 447–456.
12. CAMPBELL I. K., RICH M. J., BISCHOF R. J. and HAMILTON J. A. (2000): The colony-stimulating factors and collagen-induced arthritis: exacerbation of disease by M-CSF and G-CSF and requirement for endogenous M-CSF. *J. Leukoc. Biol.*, **68**, 144–150.
13. CECCHINI M. G., DOMINGUEZ M. G., MOCCI S., WETTERWALD A., FELIX R., FLEISCH H., CHISHOLM O., HOFSTETTER W., POLLARD J. W. and STANLEY E. R. (1994): Role of colony-stimulating factor-1 in the establishment and regulation of tissue macrophages during postnatal development of the mouse. *Development*, **120**, 1357–1372.
14. CENCI E., BARTOCCI A., PUCCETTI P., MOCCI S., STANLEY E. R.

- and BISTONI F. (1991): Macrophage colony-stimulating factor in murine candidiasis: serum and tissue levels during infection and protective effect of exogenous administration. *Infect. Immun.*, **59**, 868–872.
15. CHEERS C. and STANLEY E. R. (1988): Macrophage production during murine listeriosis: colony-stimulating factor 1 (CSF-1) and CSF-1-binding cells in genetically resistant and susceptible mice. *Infect. Immun.*, **56**, 2972–2978.
 16. CHEERS C., HILL M., HAIGH A. M. and STANLEY E. R. (1989): Stimulation of macrophage phagocytic but not bactericidal activity by colony-stimulating factor 1. *Infect. Immun.*, **57**, 1512–1516.
 17. COSTELLOE E. O., STACEY K. J., ANTALIS T. M. and HUME D. A. (1999): Regulation of the plasminogen activator inhibitor-2 (PAI-2) gene in murine macrophages. Demonstration of a novel pattern of responsiveness to bacterial endotoxin. *J. Leukoc. Biol.*, **66**, 172–182.
 18. DAI X. M., RYAN G. R., HAPPEL A. J., DOMINGUEZ M. G., RUSSELL R. G., KAPP S., SYLVESTRE V. and STANLEY E. R. (2002): Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood*, **99**, 111–120.
 19. DENIS M. (1991): Killing of *Mycobacterium tuberculosis* within human monocytes: activation by cytokines and calcitriol. *Clin. Exp. Immunol.*, **84**, 200–206.
 20. DENIS M. and GREGG E. O. (1991): Identification of cytokines which enhance (CSF-1, IL-3) or restrict (IFN- γ) growth of intramacrophage *Listeria monocytogenes*. *Immunol. Lett.*, **27**, 237–242.
 21. DEY A., SHE H., KIM L., BORUCH A., GURIS D. L., CARLBERG K., SEBTI S. M., WOODLEY D. T., IMAMOTO A. and LI W. (2000): Colony-stimulating factor-1 receptor utilizes multiple signaling pathways to induce cyclin D2 expression. *Mol. Biol. Cell*, **11**, 3835–3848.
 22. DOYLE A. G., HALLIDAY W. J., BARNETT C. J., DUNN T. L. and HUME D. A. (1992): Effect of recombinant human macrophage colony-stimulating factor 1 on immunopathology of experimental brucellosis in mice. *Infect. Immun.*, **60**, 1465–1472.
 23. ENGELHARDT R., MACKENSEN A. and GALANOS C. (1991): Phase I trial of intravenously administered endotoxin (*Salmonella abortus equi*) in cancer patients. *Cancer Res.*, **51**, 2524–2530.
 24. EVANS R., KAMDAR S. J., DUFFY T. M. and FULLER J. (1992): Synergistic interaction of bacterial lipopolysaccharide and the monocyte-macrophage colony-stimulating factor: potential quantitative and qualitative changes in macrophage-produced cytokine bioactivity. *J. Leukoc. Biol.*, **51**, 93–96.
 25. EVANS R., KAMDAR S. J., FULLER J. A. and KRUPKE D. M. (1995): The potential role of the macrophage colony-stimulating factor 1, CSF-1, in inflammatory responses: characterization of macrophage cytokine gene expression. *J. Leukoc. Biol.*, **58**, 99–107.
 26. EVANS R., SHULTZ L. D., DRANOFF G., FULLER J. A. and KAMDAR S. J. (1998): CSF-1 regulation of IL-6 gene expression by murine macrophages: a pivotal role for GM-CSF. *J. Leukoc. Biol.*, **64**, 810–816.
 27. FIRESTEIN G. S., XU W. D., TOWNSEND K., BROIDE D., ALVARO-GRACIA J., GLASEBROOK A. and ZVAIFLER N. J. (1988): Cytokines in chronic inflammatory arthritis. I. Failure to detect T cell lymphokines (interleukin 2 and interleukin 3) and presence of macrophage colony-stimulating factor (CSF-1) and a novel mast cell growth factor in rheumatoid synovitis. *J. Exp. Med.*, **168**, 1573–1586.
 28. FOWLES L. F., MARTIN M. L., NELSEN L., STACEY K. J., REDD D., CLARK Y. M., NAGAMINE Y., MCMAHON M., HUME D. A. and OSTROWSKI M. C. (1998): Persistent activation of mitogen-activated protein kinases p42 and p44 and ets-2 phosphorylation in response to colony-stimulating factor 1/c-fms signaling. *Mol. Cell Biol.*, **18**, 5148–5156.
 29. FRANCOIS B., TRIMOREAU F., VIGNON P., FIXE P., PRALORAN V. and GASTINNE H. (1997): Thrombocytopenia in the sepsis syndrome: role of hemophagocytosis and macrophage colony-stimulating factor. *Am. J. Med.*, **103**, 114–120.
 30. GALLO P., DE ROSSI A., SIVIERI S., CHIECO-BIANCHI L. and TAVOLATO B. (1994): M-CSF production by HIV-1-infected monocytes and its intrathecal synthesis. Implications for neurological HIV-1-related disease. *J. Neuroimmunol.*, **51**, 193–198.
 31. GREGORY S. H. and WING E. J. (1993): Macrophage colony-stimulating factor and the enhanced migration of monocytes are essential in primary but not secondary host defenses to *Listeria* organisms. *J. Infect. Dis.*, **168**, 934–942.
 32. GREGORY S. H., WING E. J., TWEARDY D. J., SHADDUCK R. K. and LIN H. S. (1992): Primary listerial infections are exacerbated in mice administered neutralizing antibody to macrophage colony-stimulating factor. *J. Immunol.*, **149**, 188–193.
 33. GRUBER M. F., WEIH K. A., BOONE E. J., SMITH P. D. and CLOUSE K. A. (1995): Endogenous macrophage CSF production is associated with viral replication in HIV-1-infected human monocyte-derived macrophages. *J. Immunol.*, **154**, 5528–5535.
 34. HALLET M. M., PRALORAN V., VIE H., PEYRAT M. A., WONG G., WITEK-GIANNOTTI J., SOULILLOU J. P. and MOREAU J. F. (1991): Macrophage colony-stimulating factor 1 (CSF-1) gene expression in human T-lymphocyte clones. *Blood*, **77**, 780–786.
 35. HAMILTON J. A. (1997): CSF-1 signal transduction: what is of functional significance? *Immunol. Today*, **18**, 313–317.
 36. HAMILTON J. A., WHITTY G. A., STANTON H. and MEAGER A. (1993): Effects of macrophage colony-stimulating factor on human monocytes: induction of expression of urokinase-type plasminogen activator, but not of secreted prostaglandin E₂, interleukin-6, interleukin-1, or tumor necrosis factor- α . *J. Leukoc. Biol.*, **53**, 707–714.
 37. HAMILTON J. A., WHITTY G. A., STANTON H., WOJTA J., GALLICHO M., McGRATH K. and IANCHES G. (1993): Macrophage colony-stimulating factor and granulocyte-macrophage colony-stimulating factor stimulate the synthesis of plasminogen-activator inhibitors by human monocytes. *Blood*, **82**, 3616–3621.
 38. HASHIMOTO S., YODA M., YAMADA M., YANAI N., KAWASHIMA T. and MOTOYOSHI K. (1996): Macrophage colony-stimulating factor induces interleukin-8 production in human monocytes. *Exp. Hematol.*, **24**, 123–128.
 39. HAYES M. P., WANG J. and NORCROSS M. A. (1995): Regulation of interleukin-12 expression in human monocytes: selective priming by interferon- γ of lipopolysaccharide-inducible p35 and p40 genes. *Blood*, **86**, 646–650.
 40. HO J. L., REED S. G., WICK E. A. and GIORDANO M. (1990): Granulocyte-macrophage and macrophage colony-stimulating factors activate intramacrophage killing of *Leishmania mexicana amazonensis*. *J. Infect. Dis.*, **162**, 224–230.
 41. HUME D. A., PAVLI P., DONAHUE R. E. and FIDLER I. J. (1988):

- The effect of human recombinant macrophage colony-stimulating factor 1 (CSF-1) on the murine mononuclear phagocyte system *in vivo*. *J. Immunol.*, **141**, 3405–3409.
42. ISBEL N. M., NIKOLIC-PATERSON D. J., HILL P. A., DOWLING J. and ATKINS R. C. (2001): Local macrophage proliferation correlates with increased renal M-CSF expression in human glomerulonephritis. *Nephrol. Dial. Transplant*, **16**, 1638–1647.
 43. JESSUP W., SQUIRES B., KRITHARIDES L., HUME D. A. and DEAN R. T. (1997): Effects of CSF-1 on cholesterol accumulation and efflux by macrophages. *Arterioscler. Thromb. Vasc. Biol.*, **17**, 18–25.
 44. KALTER D. C., NAKAMURA M., TURPIN J. A., BACA L. M., HOOVER D. L., DIEFFENBACH C., RALPH P., GENDELMAN H. E. and MELTZER M. S. (1991): Enhanced HIV replication in macrophage colony-stimulating factor-treated monocytes. *J. Immunol.*, **146**, 298–306.
 45. KAMDAR S. J., CHAPOVAL A. I., PHELPS J., FULLER J. A. and EVANS R. (1996): Differential sensitivity of mouse mononuclear phagocytes to CSF-1 and LPS: the potential *in vivo* relevance of enhanced IL-6 gene expression. *Cell. Immunol.*, **174**, 165–172.
 46. KAMDAR S. J., FULLER J. A., NISHIKAWA S. I. and EVANS R. (1997): Priming of mouse macrophages with the macrophage colony-stimulating factor 1 (CSF-1) induces a variety of pathways that regulate expression of the interleukin 6 (IL-6) and granulocyte-macrophage colony-stimulating factor (Csfgm) genes. *Exp. Cell Res.*, **235**, 108–116.
 47. KARBASSI A., BECKER J. M., FOSTER J. S. and MOORE R. N. (1987): Enhanced killing of *Candida albicans* by murine macrophages treated with macrophage colony-stimulating factor: evidence for augmented expression of mannose receptors. *J. Immunol.*, **139**, 417–421.
 48. KAWANAKA N., YAMAMURA M., AITA T., MORITA Y., OKAMOTO A., KAWASHIMA M., IWAHASHI M., UENO A., OHMOTO Y. and MAKINO H. (2002): CD14⁺, CD16⁺ blood monocytes and joint inflammation in rheumatoid arthritis. *Arthritis Rheum.*, **46**, 2578–2586.
 49. KAYASHIMA S., TSURU S., SHINOMIYA N., KATSURA Y., MOTYOYOSHI K., ROKUTANDA M. and NAGATA N. (1991): Effects of macrophage colony-stimulating factor on reduction of viable bacteria and survival of mice during *Listeria monocytogenes* infection: characteristics of monocyte subpopulations. *Infect. Immun.*, **59**, 4677–4680.
 50. KIM J. J., YANG J. S., LEE D. J., WILSON D. M., NOTTINGHAM L. K., MORRISON L., TSAI A., OH J., DANG K., DENTCHEV T., AGADJANYAN M. G., SIN J. I., CHALIAN A. A. and WEINER D. B. (2000): Macrophage colony-stimulating factor can modulate immune responses and attract dendritic cells *in vivo*. *Hum. Gene Ther.*, **11**, 305–321.
 51. KREMLEV S. G., CHAPOVAL A. I. and EVANS R. (1998): CSF-1 (M-CSF) enhances the inflammatory response of fibronectin-primed macrophages: pathways involved in activation of the cytokine network. *Nat. Immun.*, **16**, 228–243.
 52. KREMLEV S. G., CHAPOVAL A. I. and EVANS R. (1998): Cytokine release by macrophages after interacting with CSF-1 and extracellular matrix proteins: characteristics of a mouse model of inflammatory responses *in vitro*. *Cell. Immunol.*, **185**, 59–64.
 53. KUTZA J., CRIM L., FELDMAN S., HAYES M. P., GRUBER M., BEELER J. and CLOUSE K. A. (2000): Macrophage colony-stimulating factor antagonists inhibit replication of HIV-1 in human macrophages. *J. Immunol.*, **164**, 4955–4960.
 54. LE MEUR Y., TESCH G. H., HILL P. A., MU W., FOTI R., NIKOLIC-PATERSON D. J. and ATKINS R. C. (2002): Macrophage accumulation at a site of renal inflammation is dependent on the M-CSF/c-fms pathway. *J. Leukoc. Biol.*, **72**, 530–537.
 55. LEE M. T. and WARREN M. K. (1987): CSF-1-induced resistance to viral infection in murine macrophages. *J. Immunol.*, **138**, 3019–3022.
 56. MA X., GRI G. and TRINCHIERI G. (1996): A novel Ets-2-related nuclear factor is involved in transcriptional activation of the human interleukin-12 p40 gene promoter in response to interferon- γ and LPS stimulation of monocytic cells. *Ann. N. Y. Acad. Sci.*, **795**, 357–360.
 57. MA X., NEURATH M., GRI G. and TRINCHIERI G. (1997): Identification and characterization of a novel Ets-2-related nuclear complex implicated in the activation of the human interleukin-12 p40 gene promoter. *J. Biol. Chem.*, **272**, 10389–10395.
 58. MATSUDA M., SHIKATA K., WADA J., YAMAJI H., SHIKATA Y., DOI A., KOSAKA M., AKAGI H., MASUDA Y., OHMOTO Y. and MAKINO H. (1999): Increased urinary excretion of macrophage colony-stimulating factor (M-CSF) in patients with IgA nephropathy: tonsil stimulation enhances urinary M-CSF excretion. *Nephron*, **81**, 264–270.
 59. MATSUZAWA K., YOO Y. C., FUKUSHIMA A., YOSHIMATSU K., ARIKAWA J. and AZUMA I. (1997): Protective effect of mucosal administration of recombinant human macrophage colony-stimulating factor (rhM-CSF) on mucosal infection of Sendai virus in mice. *Vaccine*, **15**, 85–89.
 60. MOSS M. L., JIN S. L., BECHERER J. D., BICKETT D. M., BURKHART W., CHEN W. J., HASSLER D., LEESNITZER M. T., MCGEEHAN G., MILLA M., MOYER M., ROCQUE W., SEATON T., SCHOENEN F., WARNER J. and WILLARD D. (1997): Structural features and biochemical properties of TNF- α converting enzyme (TACE). *J. Neuroimmunol.*, **72**, 127–129.
 61. MOSS M. L., JIN S. L., MILLA M. E., BICKETT D. M., BURKHART W., CARTER H. L., CHEN W. J., CLAY W. C., DIDSBURY J. R., HASSLER D., HOFFMAN C. R., KOST T. A., LAMBERT M. H., LEESNITZER M. A., MCCAULEY P., MCGEEHAN G., MITCHELL J., MOYER M., PAHEL G., ROCQUE W., OVERTON L. K., SCHOENEN F., SEATON T., SU J. L., WARNER J., WILLARD D. and BECHERER J. D. (1997): Cloning of a disintegrin metalloproteinase that processes precursor tumour necrosis factor- α . *Nature*, **385**, 733–736.
 62. MUNN D. H. and ARMSTRONG E. (1993): Cytokine regulation of human monocyte differentiation *in vitro*: the tumor-cytotoxic phenotype induced by macrophage colony-stimulating factor is developmentally regulated by γ -interferon. *Cancer Res.*, **53**, 2603–2613.
 63. MUNN D. H., PRESSEY J., BEALL A. C., HUDES R. and ALDERSON M. R. (1996): Selective activation-induced apoptosis of peripheral T cells imposed by macrophages. A potential mechanism of antigen-specific peripheral lymphocyte deletion. *J. Immunol.*, **156**, 523–532.
 64. MUNN D. H., SHAFIZADEH E., ATTWOOD J. T., BONDAREV I., PASHINE A. and MELLOR A. L. (1999): Inhibition of T cell proliferation by macrophage tryptophan catabolism. *J. Exp. Med.*, **189**, 1363–1372.
 65. MUNN D. H., ZHOU M., ATTWOOD J. T., BONDAREV I., CONWAY S. J., MARSHALL B., BROWN C. and MELLOR A. L. (1998): Prevention of allogeneic fetal rejection by tryptophan catabolism. *Science*, **281**, 1191–1193.
 66. NAKOINZ I. and RALPH P. (1988): Stimulation of macrophage antibody-dependent killing of tumor targets by recombinant lymphokine factors and M-CSF. *Cell. Immunol.*, **116**, 331–340.

67. NIKOLIC-PATERSON D. J., LAN H. Y. and ATKINS R. C. (2001): Macrophages in immune renal injury. In NEILSON E. G. and COUSER W. G. (eds.): Immunologic renal disease. Raven Press, New York, 609–632.
68. OHTSUKI T., SUZU S., HATAKE K., NAGATA N., MIURA Y. and MOTOYOSHI K. (1993): A proteoglycan form of macrophage colony-stimulating factor that binds to bone-derived collagens and can be extracted from bone matrix. *Biochem. Biophys. Res. Commun.*, **190**, 215–222.
69. OZINSKY A., UNDERHILL D. M., FONTENOT J. D., HAJAR A. M., SMITH K. D., WILSON C. B., SCHROEDER L. and ADEREM A. (2000): The repertoire for pattern recognition of pathogens by the innate immune system is defined by cooperation between Toll-like receptors. *Proc. Natl. Acad. Sci. USA*, **97**, 13766–13771.
70. PHILLIPS W. A. and HAMILTON J. A. (1990): Colony-stimulating factor-1 is a negative regulator of the macrophage respiratory burst. *J. Cell. Physiol.*, **144**, 190–196.
71. PIERCE J. H., DI MARCO E., COX G. W., LOMBARDI D., RUGGIERO M., VARESI L., WANG L. M., CHOUDHURY G. G., SAKAGUCHI A. Y., DI FIORE P. P. and AARONSON S. A. (1990): Macrophage colony-stimulating factor 1 (CSF-1) induces proliferation, chemotaxis, and reversible monocytic differentiation in myeloid progenitor cells transfected with the human *c-fms*/CSF-1 receptor cDNA. *Proc. Natl. Acad. Sci. USA*, **87**, 5613–5617.
72. PRALORAN V., GASCAN H., PAPIN S., CHEVALIER S., TROSSAERT M. and BOURSIER M. C. (1990): Inducible production of macrophage colony-stimulating factor 1 (CSF-1) by malignant and normal human T cells. *Leukemia*, **4**, 411–414.
73. PRICE L. K., CHOI H. U., ROSENBERG L. and STANLEY E. R. (1992): The predominant form of secreted colony-stimulating factor-1 is a proteoglycan. *J. Biol. Chem.*, **267**, 2190–2199.
74. ROILIDES E., SEIN T., HOLMES A., CHANOCK S., BLAKE C., PIZZO P. A. and WALSH T. J. (1995): Effects of macrophage colony-stimulating factor on antifungal activity of mononuclear phagocytes against *Aspergillus fumigatus*. *J. Infect. Dis.*, **172**, 1028–1034.
75. ROTH P., BARTOCCI A. and STANLEY E. R. (1997): Lipopolysaccharide induces synthesis of mouse colony-stimulating factor 1 *in vivo*. *J. Immunol.*, **158**, 3874–3880.
76. ROTH P. and STANLEY E. R. (1992): The biology of CSF-1 and its receptor. *Curr. Top. Microbiol. Immunol.*, **181**, 141–167.
77. RYAN G. R., DAI X. M., DOMINGUEZ M. G., TONG W., CHUAN F., CHISHOLM O., RUSSELL R. G., POLLARD J. W. and STANLEY E. R. (2001): Rescue of the colony-stimulating factor 1 (CSF-1)-nullizygous mouse (*Csf1*(op)/*Csf1*(op)) phenotype with a CSF-1 transgene and identification of sites of local CSF-1 synthesis. *Blood*, **98**, 74–84.
78. SAIONJI K. and OHSAKA A. (2001): Expansion of CD4⁺CD16⁺ blood monocytes in patients with chronic renal failure undergoing dialysis: possible involvement of macrophage colony-stimulating factor. *Acta. Haematol.*, **105**, 21–26.
79. SAKURAI T., WAKIMOTO N., YAMADA M., SHIMAMURA S. and MOTOYOSHI K. (1998): Effect of macrophage colony-stimulating factor (M-CSF) on mouse immune responses *in vivo*. *Immunopharmacol. Immunotoxicol.*, **20**, 79–102.
80. SAMPSON-JOHANNES A. and CARLINO J. A. (1988): Enhancement of human monocyte tumoricidal activity by recombinant M-CSF. *J. Immunol.*, **141**, 3680–3686.
81. SEITZ M., LOETSCHER P., FEY M. F. and TOBLER A. (1994): Constitutive mRNA and protein production of macrophage colony-stimulating factor but not of other cytokines by synovial fibroblasts from rheumatoid arthritis and osteoarthritis patients. *Br. J. Rheumatol.*, **33**, 613–619.
82. SHIMA M., TEITELBAUM S. L., HOLERS V. M., RUZICKA C., OSMACK P. and ROSS F. P. (1995): Macrophage colony-stimulating factor regulates expression of the integrins $\alpha 4\beta 1$ and $\alpha 5\beta 1$ by murine bone marrow macrophages. *Proc. Natl. Acad. Sci. USA*, **92**, 5179–5183.
83. STACEY K. J., FOWLES L. F., COLMAN M. S., OSTROWSKI M. C. and HUME D. A. (1995): Regulation of urokinase-type plasminogen activator gene transcription by macrophage colony-stimulating factor. *Mol. Cell. Biol.*, **15**, 3430–3441.
84. STANLEY E. R., BERG K. L., EINSTEIN D. B., LEE P. S., PIXLEY F. J., WANG Y. and YEUNG Y. G. (1997): Biology and action of colony-stimulating factor-1. *Mol. Reprod. Dev.*, **46**, 4–10.
85. STROCKBINE L. D., COHEN J. I., FARRAH T., LYMAN S. D., WAGENER F., DUBOSE R. F., ARMITAGE R. J. and SPRIGGS M. K. (1998): The Epstein-Barr virus BARF1 gene encodes a novel, soluble colony-stimulating factor-1 receptor. *J. Virol.*, **72**, 4015–4021.
86. SWEET M. J., CAMPBELL C. C., SESTER D. P., XU D., McDONALD R. C., STACEY K. J., HUME D. A. and LIEW F. Y. (2002): Colony-stimulating factor-1 suppresses responses to CpG DNA and expression of Toll-like receptor 9 but enhances responses to lipopolysaccharide in murine macrophages. *J. Immunol.*, **168**, 392–399.
87. SZPERL M., ANSARI A. A., URBANOWSKA E., SZWECH P., KALINSKI P. and WIKTOR-JEDRZEJCZAK W. (1995): Increased resistance of CSF-1-deficient, macrophage-deficient, TNF- α -deficient, and IL-1 α -deficient op/op mice to endotoxin. *Ann. N.Y. Acad. Sci.*, **762**, 499–501.
88. TAKAHASHI M., HONG Y. M., YASUDA S., TAKANO M., KAWAI K., NAKAI S. and HIRAI Y. (1988): Macrophage colony-stimulating factor is produced by human T lymphoblastoid cell line, CEM-ON: identification by amino-terminal amino acid sequence analysis. *Biochem. Biophys. Res. Commun.*, **152**, 1401–1409.
89. TAKEDA K., KAISHO T. and AKIRA S. (2003): Toll-like receptors. *Annu. Rev. Immunol.*, **21**, 335–376.
90. TERANISHI A., AKADA S., SAITO S., HATAKE K. and MORIKAWA H. (2002): Macrophage colony-stimulating factor restored chemotherapy-induced granulocyte dysfunctions: role of IL-8 production by monocytes. *Int. Immunopharmacol.*, **2**, 83–94.
91. TILL K. J., LOPEZ A., SLUPSKY J. and CAWLEY J. C. (1993): C-fms protein expression by B-cells, with particular reference to the hairy cells of hairy-cell leukaemia. *Br. J. Haematol.*, **83**, 223–231.
92. TOJO N., ASAKURA E., KOYAMA M., TANABE T. and NAKAMURA N. (1999): Effects of macrophage colony-stimulating factor (M-CSF) on protease production from monocyte, macrophage and foam cell *in vitro*: a possible mechanism for anti-atherosclerotic effect of M-CSF. *Biochim. Biophys. Acta.*, **1452**, 275–284.
93. UTSUNOMIYA Y., OMURA K., YOKOO T., IMASAWA T., KAWAMURA T., ABE A., HIRANO K., MITARAI T., MARUYAMA N. and SAKAI O. (1996): Macrophage colony-stimulating factor (M-CSF) enhances proteinuria and recruitment of macrophages into the glomerulus in experimental murine nephritis. *Clin. Exp. Immunol.*, **106**, 286–296.
94. VASSILIADIS S. and ATHANASSAKIS I. (1994): Two novel colony-stimulating factor-1 (CSF-1) properties: it post-transcriptionally inhibits interferon-specific induction of class II antigens and reduces the risk of fetal abortion. *Cytokine*, **6**, 295–299.

95. WALKER T. M., BURGER C. J. and ELGERT K. D. (1993): Tumor growth alters macrophage responsiveness to macrophage colony-stimulating factor during reactivity against allogeneic and syngeneic MHC class II molecules. *Immunol. Invest.*, **22**, 463–476.
96. WANG J. M., GRIFFIN J. D., RAMBALDI A., CHEN Z. G. and MANTOVANI A. (1988): Induction of monocyte migration by recombinant macrophage colony-stimulating factor. *J. Immunol.*, **141**, 575–579.
97. WANG J., RODERIQUEZ G., ORAVECZ T. and NORCROSS M. A. (1998): Cytokine regulation of human immunodeficiency virus type 1 entry and replication in human monocytes/macrophages through modulation of CCR5 expression. *J. Virol.*, **72**, 7642–7647.
98. WARREN M. K. and RALPH P. (1986): Macrophage growth factor CSF-1 stimulates human monocyte production of interferon, tumor necrosis factor, and colony-stimulating activity. *J. Immunol.*, **137**, 2281–2285.
99. WEBB S. E., POLLARD J. W. and JONES G. E. (1996): Direct observation and quantification of macrophage chemoattraction to the growth factor CSF-1. *J. Cell Sci.*, **109**, 793–803.
100. WIKTOR-JEDRZEJCZAK W., DZWIGALA B., SZPERL M., MARUSZYNSKI M., URBANOWSKA E. and SZWECH P. (1996): Colony-stimulating factor 1-dependent resident macrophages play a regulatory role in fighting *Escherichia coli* fecal peritonitis. *Infect. Immun.*, **64**, 1577–1581.
101. WOHLWEND A., BELIN D. and VASSALLI J. D. (1987): Plasminogen activator-specific inhibitors in mouse macrophages: *in vivo* and *in vitro* modulation of their synthesis and secretion. *J. Immunol.*, **139**, 1278–1284.
102. ZISMAN E., WAISMAN A., BEN-YAIR E. and TARTAKOVSKY B. (1993): Production of colony-stimulating factor 1 by T cells: possible involvement in their interaction with antigen-presenting cells. *Cytokine*, **5**, 309–318.

Received in January 2003

Accepted in February 2003